Table 1. Treatment-related toxicities and survival outcomes					
Toxicity	**Group**	**Outcome**	**Survival rate (%)**	**Total n**	**p value**
Gastrointestinal toxicity (all patients)	Grade 1-2 (low-grade)	OS	74.1	44	0.260
	Grade 3-4-5 (high-grade)	OS	62.5		
Cardiac toxicity (all patients)	Present	OS	61.5	58	0.101
	Absent	OS	72.7		
	Present	DFS	38.5	53	0.177
	Absent	DFS	48.7		
Neutropenic fever (all patients)	Present	DFS	43.8	53	0.830
	Absent	DFS	47.2		
Gastrointestinal toxicity (MM only)	Grade 1-2 (low-grade)	DFS	36.8	26	0.456
	Grade 3-4-5 (high-grade)	DFS	57.1		
Mucositis (MM only)	Present	DFS	28.6	36	0.056
	Absent	DFS	57.1		
Anemia (MM only)	Grade 1-2 (low-grade)	DFS	52.6	23	0.263
	Grade 3-4-5 (high-grade)	DFS	35.7		

Treatment-related toxicities and survival outcomes. Survival rates are expressed as percentages at the end of follow-up. Total n=number of patients included in each analysis. For toxicity analyses, the highest (maximum) grade observed during conditioning and mobilization was used
MM: Multiple myeloma, OS: Overall survival, DFS: Disease-free survival